

Psychological Distress and Associated Factors among Patients with Hemophilia Living with HIV: The Japan Cohort Study of HIV-Infected Patients through Blood Products



Makiko Mieno^{1,*}, Miyuki Kawado², Shuji Hashimoto³, Kagehiro Amano⁴, Miwa Ogane⁵, Gaku Okamoto⁶, Hiroyuki Gatanaga⁵, Satoshi Higasa⁷, Hiroshi Yatsunami⁸ and Takuma Shirasaka⁶

¹Department of Medical Informatics, Center for Information, Jichi Medical University, Shimotsuke, Japan

²Department of Epidemiology and Biostatistics, National Institute of Public Health, Wako, Japan

³Fujita Health University, Toyoake, Japan

⁴Department of Laboratory Medicine, Tokyo Medical University Hospital, Tokyo, Japan

⁵AIDS Clinical Center, National Center for Global Health and Medicine, Tokyo, Japan

⁶AIDS Medical Center, National Hospital Organization Osaka National Hospital, Osaka, Japan

⁷Division of Hematology, Department of Internal Medicine, Hyogo College of Medicine, Nishinomiya, Japan

⁸Clinical Research Center, National Hospital Organization Nagasaki Medical Center, Nagasaki, Japan

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



*Address correspondence to this author at the Department of Medical Informatics, Center for Information, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke, Tochigi 329-0498, Japan; Tel: +81-285-58-7306; E-mail: mnaka@jichi.ac.jp

Cite as: Mieno M, Kawado M, Hashimoto S, Amano K, Ogane M, Okamoto G, Gatanaga H, Higasa S, Yatsunami H, Shirasaka T. Psychological Distress and Associated Factors among Patients with Hemophilia Living with HIV: The Japan Cohort Study of HIV-Infected Patients through Blood Products. Open AIDS J, 2026; 20: e18746136399483. <http://dx.doi.org/10.2174/0118746136399483260109045936>

Published: February 09, 2026



Send Orders for Reprints to reprints@benthamscience.net

Table S1. Survey items and coding.

A. Causes of worries and stress (21 items).

| | |
|----|--|
| 1 | Relationships with family |
| 2 | Relationships with people other than family |
| 3 | Love, sex |
| 4 | Marriage |
| 5 | Divorce |
| 6 | Bullying, sexual harassment |
| 7 | Purpose in life [ikigai] |
| 8 | Lack of free time |
| 9 | Income, household finances, and debt <i>etc.</i> |
| 10 | One's own illness or nursing care |
| 11 | Family member's illness or nursing care |
| 12 | Pregnancy/Childbirth |
| 13 | Childcare |
| 14 | Housework |
| 15 | Education and learning (school, studies, <i>etc.</i>) |
| 16 | Children's education |

| | |
|----|--|
| 17 | One's own work |
| 18 | Family work |
| 19 | Housing and living environment (including pollution, safety, traffic conditions) |
| 20 | Others |
| 21 | Do not know |

B. Consultation about worries and stress (15 items).

| | |
|----|---|
| 1 | Family |
| 2 | Friends or acquaintance |
| 3 | Teachers, colleagues, or supervisors |
| 4 | Consultation in public institutions |
| 5 | Consultation desk of patient organization (incl. telephone) |
| 6 | Counseling service at a private institution |
| 7 | Physician at medical institution |
| 8 | Nurse at medical institution |
| 9 | Medical social worker at medical institution |
| 10 | Clinical psychologist or counselor at medical institution |
| 11 | Counseling section of TV, radio, newspaper |
| 12 | Consultations other than 1 to 11 |
| 13 | Cannot consult anyone |
| 14 | Want to consult but do not know where to consult |
| 15 | Haven't consulted anyone because there's no need |

Note: Items are based on the Comprehensive Survey of Living Conditions in Japan (Ministry of Health, Labour and Welfare). All items were coded as 0 = No, 1 = Yes.




Fig. (S1). Forest plot of adjusted ORs for the significant stressor and consultation.